arXiv:hep-ph/9807244v1 3 Jul 1998

UAB-FT-444
July, 1998
hep-ph/9807244

# QUANTUM SUSY SIGNATURES IN LOW AND HIGH ENERGY PROCESSES

Joan SOLÀ[1]

Grup de Física Teòrica
and
Institut de Física d'Altes Energies

Universitat Autònoma de Barcelona
08193 Bellaterra (Barcelona), Catalonia, Spain

## ABSTRACT

In the search for phenomenological evidence of supersymmetry through the indirect method of quantum signatures, it is useful to seek correlations of the non-standard quantum effects in low and high energy proceses, such as those involving on one hand the properties of the $B$-mesons and on the other hand the physics of the top quark and of the Higgs bosons. There are regions of the MSSM parameter space where the potential quantum SUSY signatures in the two energy regimes are strongly interwoven and therefore the eventual detection of these correlated quantum effects would strongly point towards the existence of underlying supersymmetric dynamics.

---

[1] Invited talk at the *5th International Workshop on High Energy Physics Phenomenology* (WHEPP-5), Inter-University Centre for Astronomy and Astrophysics (IUCAA), Pune, India , January 12-26, 1998. To appear in the Proceedings.

Supersymmetry (SUSY) is perhaps the only known framework beyond the Standard Model (SM) which is capable of extending non-trivially the quantum field theoretical structure of the conventional strong and electroweak interactions while keeping all the necessary ingredients insuring internal consistency, such as gauge invariance and renormalizability. A major goal of SUSY is to produce a unified theory of all the interactions, including gravity. At present, the simplest and most popular realization of this idea, namely the Minimal Supersymmetric Standard Model (MSSM) [1], is being thoroughly scrutinized by experiment and it has successfully passed all the tests up to now. In particular, the global fit analyses to a huge number of indirect precision data within the MSSM are comparable to those in the SM [2]. One could naively think that this achievement of the MSSM is basically trivial because of the additional number of parameters involved, but this is not necessarily true on several accounts. First, the global fit analyses to a huge number of indirect precision data within the MSSM are only slightly worse than in the SM precisely because of the larger number of parameters [2]. Second, the healthy status of the MSSM from the point of view of the precision measurements is *not* shared at all by any of the known alternative extensions of the SM, like e.g. anyone of the multifarious extended Technicolour models. Third, the MSSM offers a starting point for a successful Grand Unified (GUT) framework where a radiatively stable low-energy Higgs sector can survive. Fourth, when embedding the MSSM into a GUT the running of the gauge coupling constants of the $SU(3)_c \times SU(2)_L \times U(1)_Y$ model, starting from the well-known initial conditions at the Fermi scale, turn out to match at a proton-decay-allowed GUT scale ($\sim 10^{16}\,GeV$) to within less than one standard deviation[2]. In contrast, the same running for a conventional GUT fails to unify the gauge couplings by more than $6\,\sigma$ away of a purported unifying scale which, to make things even worse, it is already ruled out by the unreported observation of proton decay. Putting things together it is well justified, we believe, to keep alive all efforts on all fronts trying to discover a supersymmetric particle. Undoubtedly, the next Tevatron run, and of course also the advent of the LHC and the possible construction of an $e^+\,e^-$ supercollider (NLC), should offer us a gold-plated scenario for finding a direct signal of SUSY. Then we should definitely see whether our faith has paid off.

In the meanwhile, and in view that supersymmetric particles must be quite heavy, one naturally looks for "quantum signatures" of the new physics by means of the indirect method of high precision measurements. In this talk we wish to stress the possibility of seeing large virtual effects of SUSY through the correlation of the quantum effects in the low and high energy domains. Thus on the one hand at high energies we have above all the physics of the top quark and the potential existence of the Higgs bosons. Indeed, one expects that a first hint of Higgs activity, if ever, should appear in concomitance with the detailed studies of top quark phenomenology. On the other hand, in the low energy domain, $B$-meson physics could be a serious competitor to high energy processes in the search for extensions of the SM. Not in vain the restrictions placed by e.g. radiative $B^0$ decays $\bar{B}^0 \to X_s\,\gamma$, i.e.

---

[2] Relative to the quantity $(\alpha_3^{-1}(M_Z) - \alpha_2^{-1}(M_Z))/(\alpha_2^{-1}(M_Z) - \alpha_1^{-1}(M_Z)) = (b_3 - b_2)/(b_2 - b_1)$ [3] which is measurable and theoretically predictible, $b_i$ being the one-loop $\beta$-function coefficients in the SUSY framework.

$$b \to s\,\gamma\,, \tag{1}$$

on the global fit analyses [2] to indirect precision electroweak data have played a significant role. Although at present the SM prediction on the decay (1) is only a little bit higher than the CLEO results [4] and the discrepancy seems no longer statistically significant, the precise knowledge of this observable is particularly useful to restrain new forms of physics beyond the SM. In fact, in the absence of SUSY, the CLEO data alone [4] on the decay (1) preclude general Type II two-Higgs-doublet models (2HDM's) [5] involving typical charged Higgs masses $M_{H^\pm} \lesssim 250\,GeV$ [6], thus barring the possibility of the non-standard top quark decay $t \to H^+\,b$. Still we should point out that the inclusion of the ALEPH data [7], with a larger central value, would not completely close this channel. In any case a bound on the charged Higgs mass is there and stems from the fact that charged Higgs bosons of $\mathcal{O}(100)\,GeV$ interfere constructively with the SM amplitude of the decay (1) and render a final value of $BR(b \to s\,\gamma)$ exceedingly high. This situation can be remedied in the MSSM, where there may be a compensating contribution from relatively light charginos and stops which tend to cancel the Higgs effects [3]. Thus, we see that from $B$-meson physics a door is still well open within the MSSM context for the non-SM top quark decay $t \to H^+\,b$ [9] which could compete with the SM mode $t \to W^+\,b$. Moreover, charged Higgs exchange may also play a role in semileptonic $B$-meson decays, especially into $\tau$-leptons,

$$b \to c\,\tau^-\,\bar{\nu}_\tau\,, \tag{2}$$

where the $2HDM$ effects could significantly modify the SM expectations [10]. Remarkably enough, the MSSM quantum corrections to $B$-meson decay processes (1)-(2) on the one hand, and to the top quark decays

$$\begin{aligned} t &\to W^+\,b\,,\\ t &\to H^+\,b\,, \end{aligned} \tag{3}$$

and the $\tau$-lepton charged Higgs decay

$$H^+ \to \tau^+\,\nu_\tau \tag{4}$$

---

[3] The recent calculation of the QCD NLO effects in the MSSM within the "heavy squark-gluino effective theory" significantly reduces the previus (LO) values of the lightest chargino-stop masses by which the SUSY contribution produces a large destructive interference with the charged-Higgs boson contribution [8]. However, the final values of the sparticle masses remain in the few hundred $GeV$ range.

on the other, can be correlated in certain regions of parameter space; e.g. they could be both maximized in the same domain. This would be a dramatic example of low and high energy correlations that could be a clue to the discovery of "virtual SUSY".

Ultimately, the potential connection between the quantum effects in the two energy regimes stems from the role played by the Yukawa sector. Within the SM the physics of the top quark is intimately connected with that of the Higgs sector through the Yukawa couplings. However, if this is true in the SM, the more it should be in the MSSM where both the Higgs and the top quark sectors are virtually "doubled" with respect to the SM. As a consequence, the Yukawa coupling sector is richer in the supersymmetric model than in the standard one. This could greatly modify the phenomenology already at the level of quantum effects on electroweak observables. As a matter of fact in the MSSM the bottom-quark Yukawa coupling may counterbalance the smallness of the bottom mass at the expense of a large value of $\tan\beta$ –the ratio $v_2/v_1$ of the vacuum expectation values of the two Higgs doublets– the upshot being that the top-quark and bottom-quark Yukawa couplings in the superpotential

$$h_t = \frac{g\,m_t}{\sqrt{2}\,M_W\,\sin\beta} \quad , \quad h_b = \frac{g\,m_b}{\sqrt{2}\,M_W\,\cos\beta}\,, \tag{5}$$

can be of the same order of magnitude, perhaps even showing up in "inverse" hierarchy: $h_t < h_b$ for $\tan\beta > m_t/m_b$. Notice that due to the perturbative bound $\tan\beta \lesssim 60$ one never reaches a situation where $h_t << h_b$. In a sense $h_t \sim h_b$ could be judged as a natural relation in the MSSM. Thus from the practical point of view, one should not dismiss the possibility that the bottom-quark Yukawa coupling could play a momentous role in the phenomenology of $B$-meson decays and top quark and MSSM Higgs boson decay and production, to the extend of drastically changing standard expectations on the observables associated to them, such as decay widths and cross-sections.

Needless to say, direct top quark decays into SUSY particles are in principle possible as well, both as 2-body and 3-body final states. Among the 2-body channels carrying an explicit SUSY signature, the following decays stand out:

$$\begin{array}{rl} \text{i)} & t \rightarrow \tilde{t}_i\,\chi^0_\alpha, \\ \text{ii)} & t \rightarrow \tilde{b}_i\,\chi^+_\alpha, \\ \text{iii)} & t \rightarrow \tilde{t}_i\,\tilde{g}\,. \end{array} \tag{6}$$

Therein, $\tilde{t}_i$, $\tilde{b}_i$, $\chi^+_i$, $\chi^0_\alpha$, $\tilde{g}$ ($i = 1, 2$; $\alpha = 1, 2, ..., 4$) denote stop, sbottom, chargino, neutralino and gluino sparticles, respectively. (Decay iii assumes that a light gluino window is still open, though it is almost ruled out at present.) Also quite a few three-body top quark decays are possible in the MSSM and have been studied in detail [11]. Here, however, we assume that sparticles (except perhaps the stop) are heavy enough that the above



Figure 1: SUSY-QCD and SUSY-EW one-loop vertices for $t \to H^+ b$ and $t \to W^+ b$. The $\Psi$'s are chargino and neutralino unphysical mass-eigenstates related to the physical mass-eigenstates ($\chi$'s) as explained in Ref. [11].

direct top SUSY decays ii)-iii) are forbidden. We are thus mainly concerned with the SM decay and the charged Higgs decay of the top quark. In fact, while the decays (3) are not necessarily MSSM processes, we wish to study whether hints of SUSY can be recognized out of them from pure supersymmetric quantum effects.

In the following we will display some numerical results accounting for the potential supersymmetric quantum corrections underlying the low and high energy decays (2)-(4) and comment on the possibility of seeing correlated SUSY quantum signatures in low and high energy processes. We perform the analysis of quantum effects in the on-shell $G_F$-scheme, which is characterized by the set of inputs $(G_F, M_W, M_Z, m_f, M_{SUSY}, ...)$, and we use the process (4) to define (and renormalize) $\tan\beta$ [9]. The supersymmetric strong (SUSY-QCD) and the supersymmetric electroweak (SUSY-EW) one-loop vertex diagrams for the charged Higgs decay of the top quark, $t \to H^+ b$, are displayed in Fig. 1. The bottom mass corrections in Fig. 2, as well as the corresponding $\tau$-lepton mass counterterms associated to our definition of $\tan\beta$, turn out to be very important. The corresponding diagrams for the standard top quark decay, $t \to W^+ b$, are obtained by just replacing $H^+$ with $W^+$ in Fig. 1, but the mass counterterms of Fig. 2 play no role since in this case there is no need to renormalize $\tan\beta$. Both decay processes (3) are well understood in the MSSM [9, 12].

Here we shall mainly report on $t \to H^+ b$ because the standard top quark decay gives a small yield. In fact, in the on-shell $G_F$-scheme the SUSY quantum corrections to the standard decay $t \to W^+ b$ are negative and of the order of a few per cent (except in some unlikely cases). Therefore, they approximately cancel out against the positive "SM" electroweak contribution [4] (of the same order of magnitude) leaving the ordinary

---

[4] Within the MSSM context, the "SM" electroweak contribution is defined to be the one obtained



Figure 2: Relevant bottom and $\tau$-lepton mass counterterms for $t \to H^+ b$ and $H^+ \to \tau^+ \nu_\tau$. Here $M_{LR}^{b,\tau} = A_{b,\tau} - \mu \tan\beta$, and $M_{LR}^t = A_t - \mu \cot\beta$, with $\mu$ the higgsino mass parameter and the $A$'s the corresponding trilinear soft SUSY-breaking couplings.

QCD effects ($\simeq -10\%$) as the net MSSM corrections (Cf. Fig. 3a). Hence no significant imprint of underlying SUSY dynamics is left behind $\Gamma(t \to W^+ b)$ and in this way we are naturally led to a closer examination of the charged Higgs decay of the top quark. To be sure, $t \to H^+ b$ has been object of many studies in the past (Cf. [9] and references therein), mainly within the context of 2HDM's, and it is being thoroughly searched in recent analyses at the Tevatron [13]. Notwithstanding, no systematic treatment of the MSSM quantum effects existed in the literature until very recently [9].

To appraise the relative importance of the various types of MSSM effects on $\Gamma(t \to H^+ b)$, in Figs. 3b-3d we provide plots for the correction to the partial width as a function of $\tan\beta$, the lightest sbottom mass ($m_{\tilde{b}_1}$) and the gluino mass ($m_{\tilde{g}}$) respectively, reflecting also the various individual contributions. Specifically, we show in these figures:

- (i) The supersymmetric electroweak contribution from genuine ($R$-odd) sparticles (denoted $\delta_{\rm SUSY-EW}$), i.e. from sfermions (squarks and sleptons), charginos and neutralinos;

- (ii) The electroweak contribution from non-supersymmetric ($R$-even) particles ($\delta_{EW}$). It is composed of two distinct types of effects, namely, those from Higgs and Goldstone bosons (collectively called "Higgs" contribution, and denoted $\delta_{\rm Higgs}$) plus the leading SM effects from conventional fermions ($\delta_{\rm SM}$):

  $$\delta_{EW} = \delta_{\rm Higgs} + \delta_{\rm SM}\,; \tag{7}$$

  The remaining non-supersymmetric electroweak effects are subleading and are neglected.

- (iii) The strong supersymmetric contribution (denoted by $\delta_{\rm SUSY-QCD}$) from squarks and gluinos;

---

after decoupling the $R$-odd effects and letting the mass of the CP-odd Higgs boson of the MSSM go to infinity [12].

- (iv) The strong contribution from conventional quarks and gluons (labelled $\delta_{\rm QCD}$); and

- (v) The total MSSM contribution, $\delta_{\rm MSSM}$, namely, the net sum of all the previous contributions:

$$\delta_{\rm MSSM}=\delta_{\rm SUSY-EW}+\delta_{EW}+\delta_{\rm SUSY-QCD}+\delta_{\rm QCD}. \tag{8}$$

We remark in Fig. 3d the local maximum in the gluino contribution (around $m_{\tilde{g}}=500\,GeV$) which entails a net MSSM correction $\delta_{\rm MSSM}\simeq 20\%$ and $75\%$ at $\tan\beta=30$ and 60 respectively. This decay is manifestly much more sensitive to heavy (not light!) gluinos ($\delta_{\rm SUSY-QCD}\simeq 0$ for $m_{\tilde{g}}=0$). Only for superheavy gluinos $m_{\tilde{g}}>>1\,TeV$ the effect eventually decouples.

We may easily understand the reason why $t\rightarrow W^+\,b$ cannot generate comparably large quantum SUSY signatures as $t\rightarrow H^+\,b$. The counterterm configuration associated to vertices involving gauge bosons and conventional fermions does *not* involve the term [14, 9] $\delta m_b/m_b$ (Cf. Fig.2) –which grows (approx.) linearly with $\tan\beta$ – so that one cannot expect similar enhancements. Therefore, the SUSY-QCD corrections to $t\rightarrow W^+\,b$ are not foreseen to be particularly significant in this case, but just of order $\alpha_s(m_t)/4\,\pi$. This is borne out by the numerical analysis in Fig.3a. The only hope for gauge boson interactions with top and bottom quarks to develop sizeable radiative corrections was to appeal to large non-oblique corrections triggered by the Yukawa terms (5). However, even in this circumstance the results are rather disappointing. For, at large $\tan\beta\geq m_t/m_b$ the bottom quark Yukawa coupling (the only relevant one in these conditions) gives a contribution of order [12]

$$\frac{\alpha_W}{4\,\pi}\,\frac{m_b^2\,\tan^2\beta}{M_W^2}\gtrsim\frac{\alpha_W}{4\,\pi}\,\frac{m_t^2}{M_W^2}\,, \tag{9}$$

which is numerically very close to the strong contribution $\alpha_s(m_t)/4\,\pi$. In contrast, the typical SUSY-QCD effect on the decay $t\rightarrow H^+\,b$ (the leading type of effect for this mode) in the limit where there is no large hierarchy between the sparticle masses is of order [9]

$$C_F\,\frac{\alpha_s}{4\,\pi}\;\tan\beta\,, \tag{10}$$

where $C_F=4/3$ is the eigenvalue of the quadratic Casimir operator of $SU(3)$ in the fundamental representation. The ratio between (10) and (9) reads

$$C_F\,\left(\frac{\alpha_s}{\alpha_W}\right)\,\left(\frac{M_W}{m_t}\right)^2\tan\beta=\mathcal{O}(1)\,\tan\beta\,. \tag{11}$$

Consequently, in the entire high $\tan\beta$ regime the SUSY-QCD effects on the $t\,b\,H^{\pm}$-vertex are expected to be a factor of order $\tan\beta$ larger than the SUSY-QCD and Yukawa coupling effects on $t\,b\,W^{\pm}$ [5].

---

[5] Of course, inclusion of the leading (Yukawa type) electroweak effects on $t\rightarrow H^+\,b$ further enhances the ratio (11). The "exact" numerical result is contained in Fig. 3b.



(a) (b)

(c) (d)

Figure 3: **(a)** The total (electroweak and strong) SUSY correction to $\Gamma(t \rightarrow W^+ b)$ for given sets of parameters. Notation as in Ref. [9]. **(b)** The SUSY-EW, SUSY-QCD, standard QCD and full MSSM corrections to $\Gamma(t \rightarrow H^+ b)$ as a function of $\tan\beta$; **(c)** As in (b), but as a function of the lightest sbottom mass; **(d)** As in (b), but as a function of the gluino mass.



(a) (b)

Figure 4: **(a)** The 95% C.L. exclusion plot in the $(\tan\beta, M_{H^\pm})$-plane for $\mu = -90\,GeV$ and remaining parameters similar to Fig. 3. Shown are the tree-level (dashed), QCD-corrected (dotted) and fully MSSM-corrected (continuous) contour lines. The excluded region in each case is the one lying below the curve; **(b)** As in (a), but for a $\mu > 0$ scenario characterized by a heavier SUSY spectrum which makes the analysis compatible with perturbation theory.

As a practical application of the previous analysis of SUSY quantum effects on $t \rightarrow H^+\,b$, let us consider the implications derived from the non-observation of an excess of $\tau$-events from $H^\pm$ decays (4) at the Tevatron [13]. Our definition of $\tan\beta$ from the vertex associated to the decay (4) allows to renormalize the $t\,b\,H^\pm$-vertex in perhaps the most convenient way to deal with $t \rightarrow H^+\,b$. Indeed, from the practical point of view, we should recall the excellent methods for $\tau$-identification developed by the Tevatron collaborations and recently used by CDF to study the permitted region in the $(\tan\beta, M_H)$-plane [13]. However, we wish to show that this analysis may undergo dramatic changes when we incorporate the MSSM quantum effects [15]. Although CDF utilizes inclusive $\tau$-lepton tagging, for our purposes it will suffice to focus on the exclusive final state $(l, \tau)$, with $l$ a light lepton, as a means for detecting an excess of $\tau$-events [16]. To be precise, we are interested in the $t\,\bar{t}$ cross-section leading to the decay sequences $t\,\bar{t} \rightarrow H^+\,b, W^-\,\bar{b}$ and $H^+ \rightarrow \tau^+\,\nu_\tau$, $W^- \rightarrow l\,\bar{\nu}_l$, and *vice versa.* The relevant quantity can be easily derived from the measured value of the canonical cross-section $\sigma_{t\bar{t}}$ for the standard channel $t \rightarrow b\,l\,\nu_l$, $\bar{t} \rightarrow b\,q\,q'$, after inserting appropriate branching fractions, namely [15]

$$\sigma_{l\tau} = \left[\frac{4}{81}\,\epsilon_1 + \frac{4}{9}\,\frac{\Gamma(t \rightarrow H\,b)}{\Gamma(t \rightarrow W\,b)}\,\epsilon_2\right]\,\sigma_{t\bar{t}}\,. \tag{12}$$

The first term in the bracket comes from the SM top quark decay, and for the second term we assume (at high $\tan\beta$) 100% branching fraction of $H^+$ into $\tau$-lepton, as explained before. Finally, $\epsilon_i$ are detector efficiency factors. Thus, in most of the phase space available for top decay the bulk of the cross-section (12) is provided by the contribution of $\Gamma(t \rightarrow H^+\,b)$. Consequently, the observable (12) should be highly sensitive to MSSM quantum effects.

In Figs. 4a and 4b we derive the (95% C.L.) excluded regions for $\mu < 0$ and $\mu > 0$, respectively. (In the $\mu > 0$ case we choose a heavier SUSY spectrum in order that the correction remains perturbative [15].) We point out that the numerical results in Figs. 4a-4b include the restrictions on the MSSM parameter space placed by the radiative $B$-meson decay, eq.(1) [6]. From inspection of these figures it can hardly be overemphasized that the MSSM quantum effects on the CDF analysis [13] can be dramatic. In particular, while for $\mu < 0$ the MSSM-corrected curve is significantly more restrictive than the QCD-corrected one, for $\mu > 0$ the bound essentially disappears from the perturbative region ($\tan\beta \lesssim 60$).

We point out the recent work of Ref. [18] claiming that the bound on the $(\tan\beta, M_H)$-plane substantially improves using Tevatron data in the $b\bar{b}\tau^+\tau^-$ channel. Nonetheless this calculation was performed by considering only conventional QCD corrections and hence it could still undergo significant MSSM radiative corrections. The potentially large effects not included in that paper stem from the production mechanism of the CP-odd Higgs boson $A^0$ (through $b\bar{b}$-fusion) before it decays into $\tau^+\tau^-$ pairs. Indeed, the $b\bar{b}\,A^0$ vertex is known [19] to develop important MSSM corrections in the relevant regions of the $(\tan\beta, M_H)$-plane purportedly "excluded" by the tree-level analysis of Ref.[18]. Therefore, a detailed re-examination of the excluded region at the quantum level is in order within the context of the MSSM before jumping into conclusions.

We complement the high energy analysis by considering the possible impact of the MSSM quantum effects on the determination of the branching ratio $BR(t \rightarrow W^+\,b)$ from the measurement of the top quark production cross-section at the Tevatron. The observed top quark production cross-section is equal to the Drell-Yan production cross-section convoluted over the parton distributions times the squared branching ratio. However, in the framework of the MSSM, we rather expect a generalization of this formula in the following way (schematically),

$$\begin{aligned} \sigma_{\rm obs.} &= \int dq\,d\bar{q}\;\sigma(q\,\bar{q} \rightarrow t\,\bar{t})\;\times |BR(t \rightarrow W^+\,b)|^2 \\ &+ \int dq\,d\bar{q}\;\sigma(q\,\bar{q} \rightarrow \tilde{b}_a\,\bar{\tilde{b}}_a)\;\times |BR(\tilde{b}_a \rightarrow \chi_1^-\,t)|^2 \times |BR(t \rightarrow W^+\,b)|^2 + ... \end{aligned} \tag{13}$$

We assume that gluinos are much heavier than squarks, so that their contribution to this cross-section through $q\,\bar{q} \rightarrow \tilde{g}\,\tilde{g}$ followed by $\tilde{g} \rightarrow t\,\tilde{t}_1$ is negligible. From eq.(13) we see that, if there are alternative (non-SM) sources of top quarks subsequently decaying into the SM final state, $W^+\,b$, then one cannot rigorously place any stringent lower bound on $BR(t \rightarrow W^+\,b)$ in the MSSM from the present FNAL data. Indeed, as we have seen above we could as well have non-SM top quark decay modes, such as e.g. $t \rightarrow \tilde{t}_a\,\chi^0_\alpha$ and $t \rightarrow H^+\,b$ that could serve, pictorially, as a "sinkhole" to compensate (at least in part) for the unseen source of extra top quarks produced at the Tevatron from sbottom pair production (Cf. eq.(13)). However, one usually assumes that $BR(t \rightarrow W^+\,b) \geq 50\%$ in

---

[6] For detailed plots of the MSSM domain allowed by $b \rightarrow s\,\gamma$, see e.g. Ref. [17].



Figure 5: Leading SUSY-EW Feynman diagrams, at one-loop order, correcting the partial width of $\tilde{b}_a \to \chi_1^- t$: **(a)** Self-energy diagrams; **(b)** Vertex contributions. Here $\Phi^0 = h^0, H^0, A^0$ are the neutral Higgs bosons of the MSSM, $H^\pm$ are the charged ones and the Goldstone bosons are denoted $G^0, G^\pm$. There are other possible diagrams, but they do not contribute in the Yukawa-coupling approximation.



(a)

(b)

Figure 6: **(a)** The SUSY-EW correction to $\tilde{b}_a \to \chi_1^- t$ as a function of $\tan\beta$. Fixed parameters are: $\mu = -90\,GeV$, $m_{\tilde{b}_1} = 300\,GeV$, $m_{\tilde{b}_2} = 350\,GeV$, $m_{\tilde{t}_1} = 200\,GeV$, $\tilde{\theta}_b = \pi/10$, $\tilde{\theta}_t = \pi/5$ and $M_H = 120\,GeV$; **(b)** As in (a), but as a function of $\mu$ and fixed $\tan\beta = 20$.

order to guarantee the (purportedly) standard top quark events observed at the Tevatron. Thus, from these considerations it is not excluded that the non-SM branching ratio of the top quark, $BR(t \rightarrow$"new"), could be as big as the SM one, i.e. $\sim 50\%$. We stress that at present one cannot exclude eq.(13) since the observed $t \rightarrow W^+ b$ final state involves missing energy, as it is also the case for the decays comprising supersymmetric particles. In particular, if $\tan\beta$ is large and there exists a relatively light chargino with a non-negligible higgsino component, the alternative mechanism suggested in eq.(13) could be a rather efficient non-SM source of top quarks that could compensate for the depletion in the SM branching ratio. From these considerations it is clear that the calculation of the MSSM quantum effects (Cf. Fig. 5) to the partial width of the decay [20, 21]

$$\tilde{b}_a \rightarrow \chi_i^- \, t \,, \tag{14}$$

could be indispensable to better assess how much the determination of the SM branching ratio $BR(t \rightarrow W^+ b)$ is affected in the MSSM context after plugging in the experimental number on the LHS of eq.(13).

In practice the computation of the loop contributions to $\tilde{b}_a \rightarrow \chi_i^- t$ requires not only a prescription to renormalize $\tan\beta$ (we use the same one as in the previous processes) but also a renormalization condition for the sbottom mixing angle, $\theta_{\tilde{b}}$, with an associated counterterm $\delta\theta_{\tilde{b}}$. In our formalism, the 3-point Green functions explicitly incorporate the mixed scalar field renormalization constants $\delta Z^{ab}$ $(a \neq b)$ and are therefore renormalized also in the $\theta_{\tilde{b}}$ parameter. The UV-divergent parts of the 3-point functions are cancelled against $\delta\theta_{\tilde{b}}$ by defining the latter as follows [21]:

$$\delta\theta_{\tilde{b}} = \frac{1}{2}\,(\delta Z^{12} - \delta Z^{21}) = \frac{1}{2}\,\frac{\Sigma^{12}(m^2_{\tilde{b}_2}) + \Sigma^{12}(m^2_{\tilde{b}_1})}{m^2_{\tilde{b}_2} - m^2_{\tilde{b}_1}} \,. \tag{15}$$

The QCD corrections (including the effect from gluinos) have been studied in Ref.[20]. Here we concentrate on the electroweak effects. The dominant part of them is expected to come from the Yukawa sector, so we just consider the electroweak effects of $\mathcal{O}(h_t^2)$ *and* $\mathcal{O}(h_b^2)$ that emerge for large values of the Yukawa couplings (5) in the region of high $\tan\beta$. The leading diagrams for the process (14) in this approximation are seen in Fig. 5. We display the evolution of the SUSY-EW effects as a function of $\tan\beta$ (Fig. 6a) and of $\mu$ (Fig. 6b). While in Fig. 6a $\mu = -90\,GeV$, we see in Fig. 6b that for $|\mu| > 120\,GeV$ the corrections can be above 20% for the lightest sbottom decay $(\tilde{b}_1)$. These corrections can reach even higher values on increasing $\tan\beta$ more and more within the perturbative region (Cf. Fig. 6a) and can have either sign depending on the stop and sbottom mixing angles[7]. Recall that the great sensitivity to large values of $\tan\beta$ was also the case for the

---

[7] However, care has to be exercized in order not to step on prohibited colour-breaking regions of the MSSM parameter space [21].



(a) (b)

(c) (d)

Figure 7: **(a)** The SUSY-QCD corrected ratio $R_{SUSY}$, eq.(16), as a function of $\tan\beta$ and for fixed $\mu < 0$. Also shown are the SM result, $R_{SM}$, (dotted band) and the Higgs-corrected result without SUSY effects, $R_H$. The shaded band is the experimental measurement at the $1\,\sigma$ level; **(b)** Similar to (a) but for two cases $\mu < 0$ and $\mu > 0$ and including the leading SUSY-EW effects; **(c)** The allowed region in the $(\tan\beta, M_H)$ plane for $\mu < 0$; **(d)** As in (c), but showing the differences of the $R_{SUSY}$-excluded area after including the leading SUSY-EW effects in that ratio.

MSSM quantum corrections to $t \to H^+ b$ (Cf. Fig. 3b). Nevertheless, in contrast to this latter decay, which was largely dominated by the bottom mass renormalization effects [9], the decay (14) receives non-negligible contributions from most of the diagrams in Fig. 5. As it is manifest in Fig. 6a, the (approximate) linear behaviour on $\tan\beta$ expected from bottom mass renormalization (Cf. Fig. 3b) becomes completely distorted by the rest of the contributions, especially in the high $\tan\beta$ end. In short, in contradistinction to the charged Higgs decay of the top quark, the final electroweak correction to the sbottom decays (14) cannot be simply ascribed to a single renormalization source but to the full Yukawa-coupling combined yield.

Moving now into the low energy domain, semileptonic $B$-meson decays can also reveal themselves as an invaluable probe for new physics. In the specific case of the inclusive semi-tauonic $B$-meson decays, $B^- \to \tau^- \bar{\nu}_\tau X$ (Cf. eq.(2)) one defines the following inclusive ratio

$$R = \frac{\Gamma(B^- \to \tau^- \, \bar{\nu}_\tau \, X)}{\Gamma(B^- \to l^- \, \bar{\nu}_l \, X)} \,, \tag{16}$$

where $l = e, \mu$ is a light lepton. In Figs.7a-7d we derive the bounds obtained on the $(\tan\beta, R)$ and $(\tan\beta, M_H)$ planes from the SUSY corrected ratio (16). The observable $R$ is sensitive to two basic parameters of generic 2HDM's, namely $\tan\beta$ and the (charged) Higgs mass, $M_H \equiv M_{H^\pm}$. As a consequence, for Type II 2HDM's (as in the MSSM case) the following upper bound at $1\,\sigma$ (resp. $2\,\sigma$) is claimed in the literature [22]:

$$\tan\beta < 0.49 \; (0.52) \; (M_H/\mathrm{GeV}) \,. \tag{17}$$

The bound (17) also hinges on the transition from the free quark model decay amplitude to the meson decay amplitude. In practice this is handled within the heavy quark expansion formalism [10]. Now, of course the point is whether that bound is significantly modified within the context of the MSSM, in particular after including the gluino mediated short-distance corrections. This SUSY-QCD analysis has been carried out in Ref. [23]. Here we improve that study by also including the leading SUSY-EW effects induced by large higgsino-squark-quark Yukawa couplings, again of the type (5). We find the following impact of the SUSY effects on the physics of the semi-tauonic inclusive $B$-meson decays within the framework of the MSSM. For $\mu > 0$, there could be no $\tan\beta - M_H$ bound at all (Fig. 7b). However, for the most likely case $\mu < 0$ (Figs. 7a-7b and 7c-7d), the SUSY effects further restrict the allowed region in the $(\tan\beta, M_H)$-plane as compared to eq.(17). The shaded region in Fig. 7c limited by the bold solid line is allowed at $2\,\sigma$ level by the ratio (16) after including SUSY-QCD effects. The narrow subarea between the thin solid lines is permitted at $1\,\sigma$ level only. The $2\,\sigma$ region allowed without including SUSY-QCD corrections is indicated by $R_H$, and the one excluded by $t \to H^+ \, b$ (without SUSY effects) is also shown. Using the present day sparticle mass limits and the LEP input data on $B$-meson decays we obtain at the $1\,\sigma$ $(2\,\sigma)$ level [23]:

$$\tan\beta < 0.40 \; (0.43) \; (M_H/\mathrm{GeV}) \,. \tag{18}$$

While in Figs. 7a and 7c we consider only the SUSY-QCD corrections to the ratio (16), in Figs. 7b and 7d we have also included the leading SUSY-EW effects, which amount to an additional 5-10% strengthening of the bound (18).

Summarizing, from the analysis of the potential SUSY quantum effects on the semi-tauonic decays and top quark decays, a correlation seems to emerge, namely at large $\tan\beta$ the SUSY imprint on the corresponding low energy and high energy processes are both maximized. The effect is fairly large, at the level of 20% or larger. For top decays into charged Higgs, it can be of order 50%. We have also found that at present the information on $(\tan\beta, M_H)$ as collected from $B$-meson decays is more restrictive than the one from top quark decays in certain regions of parameter space, specifically those which are phase-space inaccessible to top quark decay. However, in the phase-space accessible region, the data from top quark physics is more restrictive. Clearly, knowledge from both low energy and high energy data can be very useful to better pinpoint in the future the physical boundaries of the MSSM parameter space. Alternatively, if the two approaches would converge to a given portion of that parameter space, one could claim strong indirect evidence of SUSY.

Finally, we have also shown that at large $\tan\beta$ there are other SUSY decay processes and production mechanisms that could be severely influenced by the MSSM quantum corrections. These non-SM effects, in turn, could modify the determination of much more "down-to-earth" observables such as the branching ratio of the standard top quark decay. The unexpected and very much rewarding result of this is that although the SM decay $t \rightarrow W^+ b$ proves to be by itself rather insensitive to SUSY loop effects (Cf. Fig. 3a), the chain of intermediate mechanisms leading to top quark production (with the inclusion of potential SUSY sources of top quarks) could be, instead, much more sensitive to them (Cf. Fig. 6). At the end of the day it turns out that, despite the absence of noticeable SUSY quantum effects on the standard decay of the top quark, the determination of $BR(t \rightarrow W^+ b)$ from the observed cross-section could effectively incorporate a significant quantum SUSY signature!.

**Acknowledgements**: The author is grateful to the organizers of WHEPP-5 for finantial support and for the warm hospitality offered to him in IUCAA, Pune Univ., where the workshop was held. Discussions with D. P. Roy, P. Roy and M. Guchait are acknowledged. He is also thankful to Toni Coarasa and Jaume Guasch for their help in the preparation of this manuscript. This work has been partially supported by CICYT under project No. AEN93-0474.

# References

[1] J. F. Gunion, talk given at the *Int. Workshop on Quantum Effects in the MSSM*, Barcelona, September 9-13 (1997) [hep-ph/9801417], to appear in the proceedings, World Scientific 1998, Ed. J. Solà.

[2] W. Hollik, talk given at the *Int. Workshop on Quantum Effects in the MSSM*, Barcelona, September 9-13 (1997) [hep-ph/9711489], to appear in the proceedings, World Scientific 1998, Ed. J. Solà; W. de Boer, A. Dabelstein, W. Hollik, W. Mösle, U. Schwickerath, *Z. Phys.* **C 75** (1997) 627.

[3] M. Peskin, lectures given at the *European School of High-Energy Physics* [hep-ph/9705479].

[4] M.S. Alam et al. (CLEO Collab.) *Phys. Rev. Lett.* **74** (1995) 2885; S. Glenn (for the CLEO Collab.), talk at the Meeting of the American Physics Society, Ohio, March 1998.

[5] J.F. Gunion, H.E. Haber, G.L. Kane, S. Dawson, *The Higgs Hunters' Guide* (Addison-Wesley, Menlo-Park, 1990).

[6] M. Ciuchini, G. Degrassi, P. Gambino, G.F. Giudice, [hep-ph/9710335]; F. Borzumati, C. Greub, [hep-ph/9802391].

[7] P.M. Kluit, talk at the Int. Europhysics Conference, 1997.

[8] M. Ciuchini, G. Degrassi, P. Gambino, G.F. Giudice, [hep-ph/9806308].

[9] J. A. Coarasa, D. Garcia, J. Guasch, R.A. Jiménez, J. Solà, *Eur. Phys. J.* **C2** (1998) 373; J. Guasch, R.A. Jiménez, J. Solà, *Phys. Lett.* **B 360** (1995) 47.

[10] Y. Grossman, Z. Ligeti *Phys. Lett.* **B 332** (1994) 373.

[11] J. Guasch, J. Solà, *Z. Phys.* **C 74** (1997) 337.

[12] D. Garcia, W. Hollik, R.A. Jiménez, J. Solà, *Nucl. Phys.* **B 427** (1994) 53; A. Dabelstein, W. Hollik, R.A. Jiménez, C. Jünger, J. Solà, *Nucl. Phys.* **B 456** (1995) 75.

[13] B. Bevensee, talk given at the *Int. Workshop on Quantum Effects in the MSSM*, Barcelona, September 9-13 (1997), to appear in the proceedings, World Scientific 1998, Ed. J. Solà; F. Abe *et al.* (CDF Collab.), *Phys. Rev. Lett.* **79** (1997) 357.

[14] M. Carena, S. Pokorski, C.E.M. Wagner, *Nucl. Phys.* **B 426** (1994) 269; L.J. Hall, R. Rattazzi, U. Sarid, *Phys. Rev.* **D 50** (1994) 7048.

[15] J. Guasch, J. Solà, *Phys. Lett.* **B 416** (1998) 353.

[16] M. Guchait, D.P. Roy, *Phys. Rev.* **D 55** (1997) 7263.

[17] J. A. Coarasa, D. Garcia, J. Guasch, R.A. Jiménez, J. Solà, *Phys. Lett.* **B 425** (1998) 329.

[18] M. Drees, M. Guchait, P. Roy, *Phys. Rev. Lett.* **80** (1998) 2047.

[19] J.A. Coarasa, R.A. Jiménez, J. Solà, *Phys. Lett.* **B 389** (1996) 312.

[20] A. Djouadi, W. Hollik, C. Jünger, *Phys. Rev.* **D 55** (1997) 6975; S. Kraml, H. Eberl, A. Bartl, W. Majerotto, W. Porod, *Phys. Lett.* **B 386** (1996) 175.

[21] J. Guasch, W. Hollik, J. Solà, [hep-ph/9802329].

[22] Y. Grossman, H.E. Haber, Y. Nir, *Phys. Lett.* **B 357** (1995) 630.

[23] J.A. Coarasa, R.A. Jiménez, J. Solà, *Phys. Lett.* **B 406** (1997) 337.